

Interactive Effect of Background Variables and Workload Parameters on the Quality of Life among Nurses Working in Highly Complex Hospital Units: A Cross-sectional Study

MILAD ABBASI¹, SOMAYEH FARHANG DEGHAN², ROHOLLAH FALLAH MADVARI³, AHMAD MEHRI⁴, MOHAMMAD HOSSEIN EBRAHIMI⁵, MOHSEN POURSADEGHYAN⁶, HOSSEIN MARIORYAD⁷, MARYAM GHALJAH⁸

ABSTRACT

Introduction: Quality of Working Life (QWL) is a vital concept in the employees' life which can confirm the efficiency of the organisation and job satisfaction of employees.

Aim: The present study investigated the interactive effects of background variables (job and demographic characteristics) and workload parameters on the Work-Related Quality of Life (WRQoL) among nurses working in highly complex hospital units (ICU, CCU and Emergency).

Materials and Methods: This was a cross-sectional study conducted in 2017, among all male and female volunteer nurses (n=840). For this purpose, NASA-Task Load Index (NASA-TLX) and WRQoL were used. A general questionnaire was also used to collect the job and demographic information of nurses. The statistical analyses were performed through MANOVA, tests between-subject's effects, Box's M, pair-wise comparisons, Bonferroni method, and multiple regression.

Results: The findings indicated that the difference in mean score of WRQoL was statistically significant in all groups of the studied background variables except gender and Body Mass Index (BMI). The relationship between the variables of age, work experience, and the number of patients per shift and scores of work-related quality of life was statistically significant. Also, based on the results of multivariate regression analysis, the variables of overtime hours, the number of patients per shift, age, and the level of workload remained in the regression model and their coefficients of influence were estimated -43%, -23%, -19%, and -15%, respectively.

Conclusion: The results indicated the simultaneous effect of the studied variables on the nurses' WRQoL. Of these, the variables of overtime hours, the number of patients per shift, age, and workload level were finally kept in the regression model to explain the most percentage of changes in the WRQoL.

Keywords: Demographic information, Job characteristics, Work-related quality of life

INTRODUCTION

Quality of Life (QoL) is a multidimensional concept, including the functional ability, health, and psychological state of the individuals, which is influenced by many important factors such as the physical and mental status, personal belief, and social relationships [1,2]. Job has been identified as one of the affecting factor on the QoL [3]. Although job is a very important source of livelihoods and a means of acquiring social positions, it can also lead to dissatisfaction and degradation of the physical and mental forces. The workplace may be full of physical, psychological and social stimuli, each of which can be a stress factor and these stressors may have negative impact on physical well-being (individual's health and performance) and psychological well-being [4,5].

The Quality of Working Life (QWL) includes individuals' feelings about all aspects of their work, such as the financial rewards, job benefits, job security, comparing their job conditions with the employees of other organisations, working conditions, career development opportunities, decision-making power, relations with their colleagues and with the organisation [6,7].

Health system reform has been the topic of interest all over the world, especially developing countries, over the past decades. Nurses are considered to be the integral and important part in any health system [8]. Among all the human resources employed in a hospital, nursing staff is of great importance due to the provision of health care for patients and being in direct communication with patients. Delivery of the optimal nursing services is associated with the quantity and quality of nurses' work in the hospital [9]. At

present, there is a shortage of health-care workers in the health system. The widespread shortage of nurses and also the high level of turnover among nurses have become a global issue [10]. The shortage of nursing workforce reduces the positive mood of the remaining workforce and increases stress caused by the high level of workload and ultimately leads to changes in the behaviour of nurses towards their jobs. Among these changes are the low level of job satisfaction, low level of productivity, high level of cognitive failure and eventually quitting the organisation [11-14].

In the United States, between 44,000 and 98,000 people die annually in the hospitals as a result of medical errors [15]. Despite the efforts made in the fields of appropriate education and employment of nurses, we are still struggling in this area. Therefore, improving the organisational effectiveness, strengthening the organisational commitment of employees, and reducing turnover in nurses is of great importance in the health system [6].

Workload, which is defined as the total amount of work that must be done by a person or a group of people over a given time period, is high among nurses and hospital staff [16]. High workload has been reported to be as one of the main sources of occupational stress among the nurses working in the Intensive Care Units (ICUs) [17,18].

Furthermore, some issues such as getting a second job, the working unit, monthly income, age, and work experience are other factors that affect the QoL of nurses [11,19-21]; each of which has independent or interactive effects on job satisfaction and QWL.

The present study aimed to investigate the interactive effects of background variables (job and demographic characteristics) and

workload parameters on the WRQoL among 840 nurses working in highly complex hospital units including the Emergency Departments (EDs), Intensive Care Units (ICUs) and Coronary Care Units (CCUs) of the hospitals affiliated to Tehran University of Medical Sciences, Tehran, Iran.

MATERIALS AND METHODS

This was a cross-sectional study conducted in 2017 among all male and female volunteer nurses (n=840) working in hospital units including EDs, ICUs and CCUs of the hospitals affiliated to Kermanshah University of Medical Sciences. This study received ethical approval from research ethics committee of Kermanshah University of Medical Sciences and all the subjects signed an approved consent form from issued by this committee.

The NASA-Task Load Index (NASA-TLX) (Version. paper and pencil) was used to determine the workload of the subjects. The WRQoL scale was applied to determine the QWL. A general questionnaire was also used to collect the job and demographic information of nurses. The questionnaires were distributed among volunteer nurses and the method of completing the questionnaires was explained. Nurses were asked to return the completed questionnaires. The process of data collection lasted for three months from March through July 2017. A brief description of the questionnaires is presented below.

NASA-Task Load Index (NASA-TLX)

The NASA-TLX is a multi-dimensional assessment tool which evaluates the perceived workload in order to assess a task, system, or team's effectiveness or other aspects of performance [22]. The NASA-TLX originally consisted of two sections; the total workload which is divided into the six sub-scales, serving as the first part of the questionnaire:

- Mental demand
- Physical demand
- Temporal demand
- Performance
- Effort
- Frustration

There are explanations for each of these sub-scales which the subject should read before rating. They are rated for each task within a 100-points range with 5-point steps. These ratings are then combined to the Task Load Index (TLX). In second section of TLX these subscales allow subjects weighing them by pairwise comparison based on the perceived importance of those. This requires the user to choose which measurement is more relevant to work. The number of times each is chosen is the weighted score. This is multiplied by the scale score for each dimension and then divided by 15 to get a workload score from 0 to 100, which is the overall TLX [23,24]. The reliability and validity of the Persian version of this questionnaire was evaluated by Mohammadi M et al., in a study on nurses' workload ($\alpha=0.897$) [23].

Work-Related Quality of Life (WRQoL) Scale

In this study, the 23-item WRQoL scale, developed by Van Laar D et al., was applied to assess the QWL [25]. This scale uses six main components to measure the QWL of employees: career and job satisfaction, work conditions, general well-being, work-life balance, stress at work, and control at work. The questions are on 5-point Likert scale with the total score range of 0-100 in each category. The higher total scores represent the higher level of QoL. The validity and reliability of the Persian version of this scale have been confirmed by Shabaninejad H et al., [26].

STATISTICAL ANALYSIS

Multivariate Analysis Of Variance (MANOVA) was used to investigate the effect of job and demographic variables (such as gender, marital

status, working unit, number of patients per shift, overtime hours per month, second job, BMI, age, work experience, and monthly income) on the mean scores of WRQoL and workload. This test allowed examining the effect of one independent variable on more than one dependent variable.

Furthermore, to analyse the differences of the dependent variables between each of the independent variables (i.e., to examine the differences between each of the variables of the QWL and workload between each of the background variables), separate variance analyses were used as tests of between-subject's effects. It is worth noting that the hypothesis of homogeneity of variance-covariance matrix was investigated using Box's M method. To compare the effects of the estimated marginal means of the categorical variables (background variables) on each of the response variables (the mean scores of QWL and workload), the pair-wise comparison method was applied using the Bonferroni method. Moreover, after investigating the significance level of research variables on the two variables of the QWL and workload, in the next step, multiple regressions was applied to examine the factors affecting the QWL. In this study, alpha and beta were set up at 0.05 and 0.20 (a power of 0.80) respectively.

RESULTS

The sample of the present study consisted of 61.4% women and 38.6% men. The mean age and BMI of the participants was 33.36 ± 7.9 years, and 24.22 ± 3.08 kg/m² respectively. The mean QWL of subjects was obtained as 75.79 ± 13.64 and the mean workload was calculated as 69.41 ± 9.57 [Table/Fig-1].

Variable	Frequency	Percentage	
Gender	Female	516	61.4
	Male	324	38.6
Marital status	Single	438	52.1
	Married	402	47.9
Hospital unit	Emergency	216	25.7
	ICU	384	45.7
	CCU	240	28.6
Extra working hours (overtime)*	<30	180	21.4
	30-60	258	30.7
	60-90	246	29.3
	>90	156	18.6
Having a second job	Yes	294	35
	No	546	65
	Mean	SD	
Number of patients per shift	7.32	3.85	
Age (years)	33.36	7.9	
BMI (kg/m ²)	24.22	3.08	
Work experience (years)	10.53	7.33	
Monthly income (1000 Toman**)	1522.86	446.41	
Quality of working life	75.79	13.64	
Workload	69.41	9.57	

[Table/Fig-1]: Description of the background variables of the participants.

*Time worked in addition to one's normal working hours

**The Iranian toman is a superunit of the official currency of Iran, the Rial

The initial assessment of the data showed that the QWL and workload scores did not include extreme outliers. A significant correlation was found between the variables of the QWL and workload using Bartlett's test of sphericity (Chi-Square=123.35, $p<0.001$). Also, using the Box's M test, the hypothesis of equality of covariance matrices of variables of the QWL and workload was examined among the different categories of background variables (Box's M=648.96, $p<0.001$); therefore, the assumption of the homogeneity of the covariance matrices of the quantitative variables

of this study (dependent) could not be approved in the different classes of the qualitative variables (independent). The results of the tests of between-subjects effects are presented in [Table/Fig-2]. This test examines the effects of each of the dependent variables individually in the groups of independent variables.

Source	Dependent variable	Group	Mean	SD	p-value
Gender	WRQoL*	Female	75.28	0.34	108.0
		Male	76.33	0.45	
	AWWL**	Female	69.76	0.44	0.163
		Male	68.69	0.58	
Marital status	WRQoL	Single	76.65	0.38	0.001
		Married	74.96	0.36	
	AWWL	Single	66.96	0.49	<0.001
		Married	71.49	0.47	
Hospital unit	WRQoL	Emergency	72.99	0.52	<0.001
		ICU	77.26	0.36	
		CCU	77.15	0.46	
	AWWL	Emergency	71.23	0.68	<0.001
		ICU	68.97	0.47	
		CCU	67.49	0.59	
Extra working hours*	WRQoL	<30	81.89	0.67	<0.001
		60-30	80.09	0.44	
		90-60	75.22	0.46	
		>90	66.01	0.58	
	AWWL	<30	66.19	0.76	0.001
		30-60	69.08	0.59	
		60-90	69.96	0.87	
		>90	71.68	0.57	
Having a second job	WRQoL	Yes	73.79	0.44	<0.001
		No	74.82	0.31	
	AWWL	Yes	69.27	0.57	0.913
		No	69.19	0.4	
Number of patients per shift	WRQoL				<0.001
	AWWL				0.057
BMI (kg/m ²)	WRQoL				0.437
	AWWL				0.059
Age (years)	WRQoL				<0.001
	AWWL				0.001
Work experience (years)	WRQoL				<0.001
	AWWL				0.147
Monthly income (1000 Toman)	WRQoL				<0.001
	AWWL				<0.001

[Table/Fig-2]: Estimated marginal means and tests of between-subjects effects.

*Work-Related Quality of Life
 **Adaptive Weighted Workload

The results of the Pillai's Trace test showed that the variable of gender had a weak but significant effect on the QWL and workload (pillai's $F_{(2,825)}=3.15$, $p=0.043$, $\text{Partial Eta}^2=0.01$). The results of the Tests of between-subject's effects also indicated that there was no significant difference between the estimated quality of life between the men and women; as the estimated workload was not significantly different among them ($p>0.1$).


The results of paired comparisons of the estimated marginal means using the Bonferroni method showed that there was no significant difference between the men and women in terms of the QWL and workload level. However, the QWL for the single individuals was higher than that for the married ones ($p=0.001$). Workload level was found to be higher in the married people than the single ones ($p=0.001$).

Based on the results of separate analysis of variance presented in [Table/Fig-2], it was found that there is a significant difference in the QWL among different departments of the hospital ($p<0.001$). There was also a significant difference among the different departments of hospitals in terms of the workload variable ($p<0.001$).


The findings of the study showed that the age of the employees had a significant multivariate effect on the response variables (Pillai's $F_{(2,825)}=21.00$, $p<0.001$, $\text{Partial Eta}^2=0.05$). The results of the tests of inter-subject effects showed that this variable (age) had a significant effect on the QWL of employees ($p<0.001$), as its effect on the employees' workload level was also significant ($p=0.001$).

The quality of working life for the emergency department staff was significantly lower than that of ICU ($p=0.001$) and CCU ($p=0.001$) staff, while the ICU and CCU staff's QWL were approximately at the same level. Meanwhile, the workload level of the emergency department staff was significantly higher than that for the ICU staff ($p=0.019$); such a difference was also observed between the emergency department staff and CCU staff ($p=0.001$). However, no significant difference was found between the employees working in the ICUs and CCUs in terms of the workload level.

Comparison of the WRQoL and the workload mean scores of the employees in the various studied departments of the hospital in terms of gender and marital status is represented in [Table/Fig-3,4]. As it is evident, the lowest WRQoL score is associated with the female and single nurses in the emergency departments.



[Table/Fig-3]: Work-related quality of life and workload of male and female nurses in different units of the hospitals.



[Table/Fig-4]: Work-related quality of life and workload of single and married nurses in different units of the hospitals.

In order to investigate the effect of the background variables and workload on the QWL of employees, the multivariate regression analysis was applied using the Forward method. The results of the regression analysis indicated that the variables of the overtime hours, number of patients per shift, age, and the workload level would ultimately remain in the model ($F_{5,134}=79.67$, $p<0.001$), as they can account for 73.9% of the changes (Adjusted $R^2=0.739$).

Based on the results of regression analysis [Table/Fig-5], the effects of the variables of overtime hours, the number of patients per shift, and age on the QWL of employees were obtained as 43% (Beta=-0.43, $p\text{-value}<0.001$), 23% (Beta=-0.23, $p\text{-value}<0.001$), and 19% (Beta=-0.19, $p\text{-value}<0.001$), respectively. The workload level (Beta=-0.15, $p\text{-value}=0.001$) had adverse effect on the employees' quality of working life.

In order to further investigate the effect of workload on the QWL of employees, the regression of sub-scales of workload was used to evaluate its effect on the quality of employees' working life, using

Independent variable	Model coefficients				
	Regression coefficient (B)	Standard error	Standardised coefficient of regression (β)	t-value	p-value
Working overtime	-5.79	0.71	-0.43	-8.17*	<0.001
Number of patients per shift	-0.81	0.21	-0.23	-3.84*	<0.001
Age	-0.33	0.09	-0.19	-3.63*	<0.001
Workload	-0.22	0.07	-0.15	-3.25*	0.001

[Table/Fig-5]: The effect of background and workload parameters on the work-related quality of life.

*Significant at the 5% level


the Forward method. The results of this analysis showed that the physical and mental workload variables could ultimately remain in the model ($F_{3,136}=9.05$, $p<0.001$), as they can explain 14.8% of the changes (Adjusted $R_2=0.148$) [Table/Fig-6].

Independent variable	Model Coefficients				
	Regression coefficient (B)	Standard error	Standardised coefficient of regression (β)	t-value	p-value
Mental demand	-0.13	0.06	-0.19	-2.19*	0.03
Physical demand	-0.16	0.06	-0.21	-2.59*	0.011

[Table/Fig-6]: The effect of workload elements on the work-related quality of life of employees.

*Significant at the 5% level

The results of regression analysis showed that the effect of physical component of the workload on the employees' QWL was -21% (Beta=-0.16, p-value=0.011), and the effect of mental was -19% (Beta=-0.13, p-value=0.03). In other words, both affecting components of the workload variable (physical and mental) had adverse effect on the QWL of employees. This is implicitly shown in [Table/Fig-7].



[Table/Fig-7]: The relationship between the quality of working life and physical and mental subscales of workload.

DISCUSSION

The present study investigated the effects of background variables and workload parameters on the nurses' QWL. The findings indicated that in general, the WRQoL was in the lower level among the female nurses, married ones, nurses working in the emergency departments, those with an overtime of over 90 hours, and those who had a second job compared to the effects of other studied variables; that is to say that the difference in mean score of WRQoL was statistically significant in all groups in terms of the studied background variables except gender and BMI. The relationship

between the variables of age, work experience, and the number of patients per shift and nurses' quality of life scores was statistically significant. Also, based on the results of multivariate regression analysis, the variables of overtime hours, the number of patients per shift, age, and the level of workload remained in the regression model and their coefficients of influence were estimated -43%, -23%, -19%, and -15%, respectively.

Although the role of gender on the nurses' WRQoL score was not statistically significant, the QoL score of female nurses was lower than that of men. Other studies investigated gender differences in the QWL also suggested that men are in more privileged positions in work environments, while women's commitment to their work is higher [27,28]. Gender differences have been proven in many of the indicators of work quality including the independence, job satisfaction, supervisor support, job security, and balance between work and life. According to previous studies, men are more likely to be paid better than women, get better organisational posts, and have more chance of flexible using of their work time and making decision on their work plan; while women spend more time on working and are more loyal to the organisation than men [27,28]. Female workload score was also higher than of men, although this difference was not statistically significant. Although both female and male nurses experienced a high level of workload, all data showed a more favourable situation for men compared to women. Women are more stressed due to unpaid home-task and have more responsibilities to their family. Studies have shown that women in the workplace have a higher level of norepinephrine than men, reflecting their high volume of workload [29,30].

The results of Pillai's trace test showed that married nurses had significantly lower QoL than unmarried ones, which could be affected by their higher level of workload resulting in a lower level of job satisfaction. This can be due to many reasons including the current situation of economic pressures in Iran, life responsibilities, and family problems for married people [31], although identifying the root causes of this issue requires more comprehensive studies. Han KT et al., examined the relationship between the QoL and marital status among the cancer patients and supervisors [32]. They observed that the QoL score was higher in single people. However, there are some studies that consider the QoL of married people higher than that of in single people [33,34], this may be due to the more emotional and social support that married people receive from their spouses.

Emergency departments of hospitals are among the most stressful working environments. This can be justified with the difficult work conditions in EDs, including the large number of patients, the need for rapid decision-making, the high volume of workload and psychological pressures, lack of resources (human and materials), and poor support. As a result, the prevalence of burnout among the nurses and doctors working in this department is usually at the moderate to high level [35,36]. Similar to the results of this study, other studies have also shown that the ED staff have lower level of QoL and more workload level [35,37].

Based on the results of some previous studies, long working hours can be associated with the higher workloads, poorer lifestyles, higher levels of stress, higher fatigue, and lower level of QoL [38,39]. Long working hours have a particularly negative effect on the interaction between work and home of employees [40]. Similarly, in the present study, with the increase in the overtime hours, the level of WRQoL decreased and the level of workload increased in nurses. Long working hours has significant effects on lifestyle, including the pattern of sleep, the discipline of everyday life, and meals; and it has been shown that people with longer working hours experience more stress [40]. Consistent with this finding, the results of tests of between-subjects effects showed that nurses without a second job have a higher level of QoL than nurses with a second job.

The results of tests of between-subjects effects showed that the variables of age and work experience of employees have a significant effect on the QWL of employees, so that by an increase in the age and work experience, the WRQoL decreases. These two variables (age and work experience) had a reverse effect on all components of the WRQoL variable. Many studies have shown that mental and physical abilities and capacities of nurses reduce with the increase in their age [41,42]; as a result, this variable, with the effect on job satisfaction, increased work stress, reduced control on work, increased exhausting effort, and frustration, can reduce the WRQoL in this high-stressed job [43].

The QWL and work-related consequences, such as job satisfaction and organisational commitment are among the important issues in the development of human and organisational resources [44].

LIMITATION

Despite its large sample size, this study had some limitations, including the cross-sectional nature of the study, the complexity of the considered topic, the presence of many influential factors on the level of WRQoL (such as the personal life, hospital laws and policies with regard to the compensation and benefits, interaction with colleagues and supervisors, work environment and working conditions, and the possibility of job growth), and using a questionnaire as a subjective measurement tool (Differences in understanding and interpretation, dishonesty of respondents).

CONCLUSION

The results indicated the simultaneous effect of the studied variables on the nurses' WRQoL; among these, the variables of overtime hours, the number of patients per shift, age and workload level was found to affect the work-related quality of life.

Funding: This article is extracted from research project No.96023. The authors would like to acknowledge the financial support of Kermanshah University of Medical Sciences for this research.

REFERENCES

- [1] Dumuid D, Olds T, Lewis LK, Martin-Fernández JA, Katzmarzyk PT, Barreira T, et al. Health-related quality of life and lifestyle behavior clusters in school-aged children from 12 countries. *The Journal of Pediatrics*. 2017;183:178-83. e2.
- [2] Polonsky WH. Understanding and assessing diabetes-specific quality of life. *Diabetes Spectrum*. 2000;13(1):36.
- [3] Islam MMB. Factors affecting quality of work life: an analysis on employees of private limited companies in Bangladesh. *Global Journal of Management and Business Research*. 2012;12(18).
- [4] Attell BK, Brown KK, Treiber LA. Workplace bullying, perceived job stressors, and psychological distress: Gender and race differences in the stress process. *Social Science Research*. 2017;65:210-21.
- [5] Malekpour F, Mohammadian Y, Mohamadpour Y, Fazli B, Hassanloei B. Assessment of relationship between quality of life and mental workload among nurses of urmia medical science university hospitals. *The Journal of Urmia Nursing and Midwifery Faculty*. 2014;12(6):499-504 [in Persian].
- [6] Lin S, Chaiear N, Khiewyoo J, Wu B, Johns NP. Preliminary psychometric properties of the Chinese version of the work-related quality of life scale-2 in the nursing profession. *Safety and Health at Work*. 2013;4(1):37-45.
- [7] Garg C, Munjal N, Bansal P, Singhal AK. Quality of work life: An overview. *International Journal of Physical and Social Sciences*. 2012;2(3):231-42.
- [8] Dai HD, Tang FI, Chen IJ, Yu S. Taiwanese version of the work-related quality of life scale for nurses: Translation and validation. *Journal of Nursing Research*. 2016;24(1):58-67.
- [9] Safari S, Mohammadi-Bolbanabad H, Kazemi M. Evaluation mental work load in nursing critical care unit with National Aeronautics and Space Administration Task Load Index (NASA-TLX). *J Health Sys Res*. 2013;9(6):613-19.
- [10] Lu H, While AE, Barrball KL. Job satisfaction and its related factors: A questionnaire survey of hospital nurses in Mainland China. *International Journal of Nursing Studies*. 2007;44(4):574-88.
- [11] Almkali MJ, FitzGerald G, Clark M. The relationship between quality of work life and turnover intention of primary health care nurses in Saudi Arabia. *BMC Health Serv Res*. 2012;12(1):314.
- [12] Portoghese I, Galletta M, Coppola RC, Finco G, Campagna M. Burnout and workload among health care workers: the moderating role of job control. *Safety and Health at Work*. 2014;5(3):152-57.
- [13] Abbasi M, Zakerian A, Mehri A, Poursadeghiyan M, Dinarvand N, Akbarzadeh A, et al. Investigation into effects of work-related quality of life and some related factors on cognitive failures among nurses. *Int J Occup Saf Ergon*. 2017;23(3):386-92.
- [14] Poursadeghiyan M, Abbasi M, Mehri A, Hami M, Raei M, Ebrahim M. Relationship between job stress and anxiety, depression and job satisfaction in nurses in Iran. *Social Sciences (Pakistan)*. 2016;11(9):2349-55.
- [15] Young G, Zavelina L, Hooper V. Assessment of workload using NASA Task Load Index in perianesthesia nursing. *Journal of Peri Anesthesia Nursing*. 2008;23(2):102-10.
- [16] Gurses AP, Carayon P, Wall M. Impact of performance obstacles on intensive care nurses' workload, perceived quality and safety of care, and quality of working life. *Health Services Research*. 2009;44(2p1):422-43.
- [17] Baggs JG, Schmitt MH, Mushlin AI, Mitchell PH, Eldredge DH, Oakes D, et al. Association between nurse-physician collaboration and patient outcomes in three intensive care units. *Critical Care Medicine*. 1999;27(9):1991-98.
- [18] Embriaco N, Papazian L, Kentish-Barnes N, Pochard F, Azoulay E. Burnout syndrome among critical care healthcare workers. *Curr Opin Crit Care*. 2007;13(5):482-88.
- [19] Hamaideh SH. Occupational stress, social support, and quality of life among Jordanian mental health nurses. *Issues in Mental Health Nursing*. 2011;33(1):15-23.
- [20] Mor Barak ME, Nissly JA, Levin A. Antecedents to retention and turnover among child welfare, social work, and other human service employees: What can we learn from past research? A review and metanalysis. *Social Service Review*. 2001;75(4):625-61.
- [21] Khammar A, Moghimian M, Ebrahimi MH, Abbasi M, Baneshi MM, Yari AR, et al. Effects of bright light shock on sleepiness and adaptation among night workers of a hospital in Iran. *Annals of Tropical Medicine and Public Health*. 2017;10(3):595.
- [22] Hart SG, Staveland LE. Development of NASA-TLX (Task Load Index): Results of empirical and theoretical research. *Advances in psychology*. 52: Elsevier; 1988. Pp. 139-83.
- [23] Mohammadi M, Mazloumi A, Zeraati H. Designing questionnaire of assessing mental workload and determine its validity and reliability among ICUs nurses in one of the TUMS's hospitals. *Journal of School of Public Health and Institute of Public Health Research*. 2013;11(2):87-96.
- [24] Hart SG, editor NASA-task load index (NASA-TLX); 20 years later. Proceedings of the human factors and ergonomics society annual meeting. Los Angeles, CA: Sage Publications Sage CA; 2006.
- [25] Van Laar D, Edwards JA, Easton S. The Work-Related Quality of Life scale for healthcare workers. *Journal of Advanced Nursing*. 2007;60(3):325-33.
- [26] Shabaninejad H, Arab M, Rashidian A, Zeraati H, Bahrami S. Quality of working life of Family Physicians in Mazandaran. *Hakim Research Journal*. 2012;15(2):178-84.
- [27] Den Dulk L, Bäck-Wiklund M, Lewis S, Redai D. Quality of life and work in a changing Europe: a theoretical framework. *Quality of Life and Work in Europe*: Springer; 2011. Pp. 17-31.
- [28] van der Lippe T, den Dulk L, van Doorne-Huiskes A, Schippers J, Lane L, Bäck-Wiklund M. Quality of life in a changing Europe. Final Report of FP6 project, [Online]. Universiteit Utrecht. Available at: [projectquality.org/files/Final% 20 Report_2009.pdf](http://projectquality.org/files/Final%20Report_2009.pdf) (accessed 9 May 2012); 2009.
- [29] Eek F, Axmon A. Gender inequality at home is associated with poorer health for women. *Scandinavian Journal of Social Medicine*. 2015;43(2):176-82.
- [30] Lundberg U, Mårdberg B, Frankenhaeuser M. The total workload of male and female white collar workers as related to age, occupational level, and number of children. *Scandinavian Journal of Psychology*. 1994;35(4):315-27.
- [31] Mehr RK, Mohammadi A, Pour ZA. Married' students attitude towards marriage: a qualitative study. *Iranian Journal of Psychiatric Nursing*. 2016;4(2):19-27.
- [32] Han KT, Kim SJ, Song H, Chun SY, Kim CO, Kim JS, et al. Associations between quality of life and marital status in cancer patients and survivors. *Asian Pacific Journal of Cancer Prevention*. 2014;15(13):5287-91.
- [33] Luttik ML, Jaarsma T, Veeger N, van Veldhuisen DJ. Marital status, quality of life, and clinical outcome in patients with heart failure. *Heart & Lung: The Journal of Acute and Critical Care*. 2006;35(1):3-8.
- [34] Wang FL, Gu XM, Hao BY, Wang S, Chen ZJ, Ding CY. Influence of marital status on the quality of life of Chinese adult patients with epilepsy. *Chinese Medical Journal*. 2017;130(1):83.
- [35] Bragard I, Fleet R, Etienne AM, Archambault P, Légaré F, Chauny JM, et al. Quality of work life of rural emergency department nurses and physicians: a pilot study. *BMC Research Notes*. 2015;8(1):116.
- [36] Potter C. To what extent do nurses and physicians working within the emergency department experience burnout: A review of the literature. *Australasian Emergency Nursing Journal*. 2006;9(2):57-64.
- [37] Rossetti AC, Gaidzinski RR, Fugulin FMT. Nursing workload in the emergency department: A methodological proposal. *Revista Latino-Americana De Enfermagem*. 2013;21(SPE):225-32.
- [38] Van der Hulst M. Long workhours and health. *Scandinavian Journal of Work, Environment & Health*. 2003;29(3):171-88.
- [39] Wu HC, Wang M-JJ. Relationship between maximum acceptable work time and physical workload. *Ergonomics*. 2002;45(4):280-89.
- [40] Maruyama S, Morimoto K. Effects of long workhours on life-style, stress and quality of life among intermediate Japanese managers. *Scandinavian Journal of Work, Environment & Health*. 1996;22(5):353-59.
- [41] Chiu M-C, Wang M-JJ, Lu C-W, Pan S-M, Kumashiro M, Ilmarinen J. Evaluating work ability and quality of life for clinical nurses in Taiwan. *Nursing Outlook*. 2007;55(6):318-26.

- [42] Golubic R, Milosevic M, Knezevic B, Mustajbegovic J. Work-related stress, education and work ability among hospital nurses. *Journal of Advanced Nursing*. 2009;65(10):2056-66.
- [43] Abbasi M, Zakerian A, Akbarzade A, Dinarvand N, Ghaljahi M, Poursadeghiyan M, et al. Investigation of the relationship between work ability and work-related quality of life in nurses. *Iranian Journal of Public Health*. 2017;46(10):1404.
- [44] Lai SL, Chang J, Hsu LY. Does effect of workload on quality of work life vary with generations? *Asia Pacific Management Review*. 2012;17(4):437-51.

PARTICULARS OF CONTRIBUTORS:

1. Research Center for Environmental Determinants of Health (RCEDH), Kermanshah University of Medical Sciences, Kermanshah, Iran.
PhD Student, Students' Scientific Research Center, Tehran University of Medical Sciences, Tehran, Iran.
2. Assistant Professor, Department of Occupational Health, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran.
3. PhD Candidate, Students Research Committee, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
4. Student, Department of Occupational Health Engineering, School of Public Health, Iranshahr University of Medical Sciences, Iranshahr, Iran.
5. Associate Professor, Environmental and Occupational Health Research Center, Shahroud University of Medical Sciences, Shahroud, Iran.
6. Assistant Professor, Health in Emergency and Disaster Research Center, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran.
7. Social Determinants of Health Research Center, Yasuj University of Medical Sciences, Yasuj, Iran.
8. Instructor, Department of Occupational Health Engineering, School of Public Health, Zabol University of Medical Sciences, Zabol, Iran.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Maryam Ghaljahi,
Instructor, Department of Occupational Health Engineering, School of Public Health,
Zabol University of Medical Sciences, Zabol, Iran.
E-mail: occmgh@gmail.com; milad8285@gmail.com

Date of Submission: **Jul 18, 2018**

Date of Peer Review: **Aug 20, 2018**

Date of Acceptance: **Sep 13, 2018**

Date of Publishing: **Jan 01, 2019**

FINANCIAL OR OTHER COMPETING INTERESTS: As declared above.