



# Effect of *Artemisia vulgaris* Hydro-Alcoholic Extract on Oxidative Stress and Inflammatory Damages in a Rat Model of Experimental Colitis

Arezoo Moini Jazani <sup>1</sup>, Sahar Shafiei <sup>2</sup>, Hosna Khazaei <sup>3</sup>, Mohammad Hashemnia <sup>4</sup>, Sajad Fakhri <sup>3</sup> and Mohammad Hosein Farzaei <sup>1,3,\*</sup>

<sup>1</sup>Traditional Medicine and Hydrotherapy Research Center, Ardabil University of Medical Sciences, Ardabil, Iran

<sup>2</sup>Student Research Committee, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, Iran

<sup>3</sup>Pharmaceutical Sciences Research Center, Health Institute, Kermanshah University of Medical Sciences, Kermanshah, Iran

<sup>4</sup>Department of Pathobiology, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

\*Corresponding author: Traditional Medicine and Hydrotherapy Research Center, Ardabil University of Medical Sciences, Ardabil, Iran. Email: mh.farzaei@gmail.com

Received 2023 August 01; Revised 2023 September 29; Accepted 2023 October 15.

## Abstract

**Background:** Herbal medicines can be used as a possible therapeutic agent in inflammatory diseases.

**Objectives:** This study aimed to determine the effect of *Artemisia vulgaris* (AV) hydro-alcoholic extract on oxidative stress and inflammatory damage in an experimental rat model of colitis.

**Methods:** Thirty male Sprague-Dawley rats were randomly divided into 6 groups: Control, colitis, sulfasalazine, AV50, AV100, and AV200. The animals of the AV groups were treated with the hydro-alcoholic extract of *A. vulgaris* via gavage for 72 h. Body weight was measured at the beginning and end of the experiment. In the end, serum levels of malondialdehyde (MDA) as a lipid peroxidation marker, antioxidants such as glutathione (GSH), superoxide dismutase (SOD) activity, serum tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), and nitric oxide (NO) levels as inflammatory biomarkers were measured. Macroscopic and microscopic damage in the rat's colon was also examined in histological studies.

**Results:** The *A. vulgaris* extract treatment dose-dependently improved colonic injury ( $P < 0.001$  to  $P < 0.05$ ) and body weight ( $P < 0.001$ ) in colitis rats. Moreover, it enhanced SOD activity and GSH levels ( $P < 0.001$ ) and reduced serum MDA, TNF- $\alpha$ , and NO levels ( $P < 0.001$  to  $P < 0.01$ ) in the rats with colitis.

**Conclusions:** Treatment with *A. vulgaris* could mitigate ulcerative colitis (UC) symptoms, which is probably attributed to its antioxidant and anti-inflammatory properties.

**Keywords:** *Artemisia vulgaris*, Ulcerative Colitis, Oxidative Stress, Inflammation, Rat

## 1. Background

Ulcerative colitis (UC) is a type of inflammatory bowel disease (IBD) that results in inflammation and ulceration of the intestinal mucosa. It is chronic, recurrent, and progressive, and its main clinical hallmarks include mucous diarrhea with blood, weight loss, abdominal pain, anemia, and fatigue (1-3). Based on a recent systematic review by Ng et al., the highest prevalence rates of UC were in North America (0.29%) and Europe (0.51%) (4). In Iran, the incidence of UC has increased over the past two decades (5, 6). The etiology of the disease is complex and unclear. However, epidemiological studies found that environmental and genetic predisposition plays a key role in inflammation-related diseases. The pathological features include immunologic abnormalities

(7), abnormal microflora of the GI (8), oxidant/antioxidant imbalance (9), elevated pro-inflammatory cytokines, and defects in mucosal integrity (10, 11).

Emerging evidence indicates that oxidative stress plays a crucial role in the development of intestinal inflammation via underlying mechanisms, including overproduction of reactive oxygen species (ROS), infiltration of immune cells, and upregulation of inflammatory cytokines (12, 13). Furthermore, many studies have shown a positive correlation between high concentrations of nitric oxide (NO) in the serum and intestinal mucosa and increased levels of pro-inflammatory cytokines in the induction of chronic UC (14, 15). Currently, immunosuppressive agents, anti-inflammatory drugs, corticosteroids,

aminosalicylates, and probiotics are commonly used to control inflammation and alleviate the risk of recurrence. However, the response rate to available drugs is poor, and there are various side effects. Therefore, it is significant to search for potential therapeutic strategies for UC. Medicinal plants are worthy and effective in treating diseases and are the source of most newly discovered drugs (16). The use of natural products to treat various gastrointestinal diseases offers an alternative therapy.

*Artemisia vulgaris* (known as mugwort) is a well-known genus of the Asteraceae family, with many species found in Iran, most of which are aromatic. This genus is native to the temperate regions of Europe, Asia, North Africa, and Africa and contains saponins, alkaloids, essential oils, phenolic acids, coumarins, and flavonoids (17, 18). Further, *A. vulgaris* called the "mother of herbs," is used as an anti-inflammatory, antioxidant, and immunomodulatory agent (19). It has also been used as an analgesic and anti-ulcerogenic (20, 21). Researchers have confirmed that *A. vulgaris* exhibits antibacterial, antiseptic, anticancer, and hepatoprotective properties, as well as therapeutic effects in metabolic disorders such as diabetes (18, 22). In this regard, recent studies have demonstrated that *A. vulgaris* can attenuate inflammatory mediators, such as cyclooxygenase-2 (COX-2) and inducible nitric oxide synthase (iNOS) (23, 24), increase antioxidant activity by upregulating nuclear factor erythroid 2-related factor 2 (Nrf2) (25), and improve damage levels in an experimental model of colitis.

## 2. Objectives

Considering the medicinally useful properties of the plant and due to the limited number of studies on its anti-ulceration effects in rats, this study investigated the efficacy of *A. vulgaris* hydroalcoholic extract on oxidative stress and inflammatory damage in an experimental model of acetic acid-induced colitis.

## 3. Methods

### 3.1. Animals

Male Sprague-Dawley rats (8 - 10 weeks old; 180 - 200 g) were purchased from the animal laboratory of the Faculty of Pharmacy, Kermanshah University of Medical Sciences (Iran). The rats were housed in a temperature-controlled room ( $22 \pm 2^\circ\text{C}$ ) with a 12-h light/dark cycle (lights on at 7:00 AM) and food and tap water ad libitum throughout the experiments. All the procedures were conducted in compliance with the National Institutes of Health guidelines for the care and use of laboratory animals (NIH;

publication No. 82-23, revised 1985 and again implemented in 1996) and approved by the Ethics Committee of Ardabil University of Medical Sciences (IR.ARUMS.REC.1400.202).

### 3.2. Preparation of *A. vulgaris* Extract

*A. vulgaris* was collected in May 2021 from Kermanshah province in western Iran and identified by the herbarium of the Department of Pharmacology, Kermanshah University of Medical Sciences, Kermanshah (voucher No. 075-014-034). Briefly, the aerial parts of the plant were powdered, extracted with 70% ethanol using the maceration technique, and filtered. The resulting dried extract was using a centrifugal evaporator (Heidolph-Laborota 4001, Germany) and stored at  $4^\circ\text{C}$  for further experiments (26).

### 3.3. Induction of Ulcerative Colitis

After 24 h of fasting, the animals were anesthetized with ketamine/xylazine (40/10 mg/kg, IM). Then, an acute UC model was induced by intrarectal administration of acetic acid (AA, 2 mL, 3% v/v, via the medical-grade polyurethane catheter, Sigma-Aldrich). After the AA injection, the rats were physically monitored to check for weight loss, diarrhea, and anorectal bleeding, which indicated the induction of colitis (27, 28).

### 3.4. Experimental Groups

The rats were randomly assigned to 6 groups ( $n = 5$  rats/group): (1) control group: Healthy rats that received 1 mL of normal saline orally (gavage); (2) colitis group: Colitis was induced with acetic acid intrarectally (2 mL of 3% in 0.9% NaCl) and the rats received 1 mL of normal saline orally; (3) sulfasalazine group: Ulcerative colitis-induced rats that received sulfasalazine as a reference drug [1 mg/kg/day; intraperitoneally (i.p.)]; (4) AV50 group: Ulcerative colitis-induced rats that received 50 mg/kg of *A. vulgaris* orally; (5) AV100 group: Ulcerative colitis-induced rats that received 100 mg/kg of *A. vulgaris* orally; (6) AV200 group: Ulcerative colitis-induced rats that received 200 mg/kg of *A. vulgaris* orally. The treatment period was 7 days before induction and 3 days after. The doses of *A. vulgaris* extracts and the duration of treatment were selected according to previous studies (29-31). The body weight of the rats was measured at the beginning and end of the interventions using an FEW (Japan) scale with a sensitivity of 1 g.

### 3.5. Collection of Samples

On the last day of the experiment, the rats were sacrificed under deep anesthesia with ketamine and xylazine (100 and 10 mg/kg i.p., respectively). Then, trunk blood was collected into a microtube, centrifuged at 3000 rpm for 10 min, and kept at -80°C for evaluation of the pro-inflammatory mediator and oxidative stress markers. In addition, the colon tissue was immediately removed and fixed in a formalin solution for macroscopic and histopathological studies.

### 3.6. Measurement of Serum NO and Tumor Necrosis Factor- $\alpha$ Levels

Serum NO and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) levels were measured using a rat ELISA kit (ZellBio GmbH, Germany) according to the manufacturer's protocols and expressed as  $\mu$ M and Pg/mL, respectively.

### 3.7. Measurement of Serum Oxidative Stress Markers

Levels of serum malondialdehyde (MDA), glutathione (GSH), and superoxide dismutase (SOD) activity were determined using a colorimetric kit (ZellBio GmbH, Germany) as per the manufacturer's instructions and expressed as nmol/mL and U/mL, respectively.

### 3.8. Macroscopic Assessment of Colonic Damage

The macroscopically visible damage of the colon was examined under a stereomicroscope and scored as described by Farzaei et al. (28). Mucosal edema, thickening, hemorrhage, hyperemia, erosions, shortening, and necrosis were assessed based on the Gerald scoring system (32).

### 3.9. Histopathological Examination of Colonic Damage

Colonic tissue samples were collected for microscopic (histopathological) evaluation and fixed in 10% buffered neutral formalin. The segments were then processed, embedded in paraffin blocks, sectioned (5 - 7  $\mu$ m) with a rotary microtome, and counterstained with hematoxylin-eosin (H&E) for analysis under a light microscope. The description and scoring of lesions were conducted based on El-Akabawy and El-Sherif (33) as follows: Morpho-architectural distortion of crypts, submucosal edema, ulceration, cryptitis, and glandular atrophy: Absent (score 0), mild (score 1,  $\leq$  10%), moderate (score 2, 10 - 15%), and intense (score 3,  $>$  50%). The final scores were calculated by summing the scores for each sample. The degree of inflammatory cell infiltration was also scored according to the following scale: 0, normal; 1, presence of inflammatory cells confined to the mucosa; 2,

present in both mucosa and submucosa; and 3, infiltrate extended into the traversal of the entire length of the colonic wall.

### 3.10. Statistical Analysis

The data were analyzed using GraphPad Prism v. 8. One-way analysis of variance (ANOVA; considering the drug as the independent factor) and two-factor mixed-model ANOVA (drug as the independent factor and time as the repeated factor), followed by Tukey's post-hoc test for multiple comparisons, were used. The results are expressed as mean + standard error of the mean (SEM), and P-values below 0.05 were considered significant.


## 4. Results

### 4.1. *A. vulgaris* Increased Body Weight in the Rat Model of Colitis

Two-way repeated-measures ANOVA showed the significant effect of time [F (1, 48) = 78.77, P < 0.0001] and treatment [F (5, 48) = 10.89, P < 0.0001] and the significant effect of time  $\times$  treatment interaction [F (5, 48) = 9.202, P < 0.0001] on the body weight of the experimental groups. Based on Figure 1, the body weights of the animals were not significant in any assigned groups on day 1 of the experiment. On the other hand, colitis rats showed significantly decreased body weights compared to control groups on day 7 of the study (P < 0.001). However, *A. vulgaris* treatment for 3 days at a 200 mg/kg concentration caused a significant increase (P < 0.001) in the body weight of the animals compared to the rat model of colitis at the end of the study (P < 0.001) (Figure 1).

### 4.2. *A. vulgaris* Diminished Serum NO and TNF- $\alpha$ Levels in the Rat Model of Colitis

As shown in Figure 2A, B, a significant difference was shown by one-way ANOVA in serum NO [F (5, 24) = 352.9, P < 0.0001] and TNF- $\alpha$  levels [F (5, 24) = 63.58, P < 0.0001] between the study groups. The post-hoc analysis indicated that serum levels of NO and TNF- $\alpha$  significantly increased in the colitis group when compared with the control group (P < 0.001). At all doses tested, treatment with *A. vulgaris* resulted in a significant decrease in serum NO and TNF- $\alpha$  levels (except in the AV50 group) in colitis rats compared to untreated colitis rats (P < 0.001) (Figure 2A, B). In addition, there was a significant difference between the sulfasalazine group and the colitis group (P < 0.001) (Figure 2A, B).



**Figure 1.** Effect of *A. vulgaris* extract treatment on body weight in acetic acid (AA)-induced colitis rats. Data are represented as mean  $\pm$  standard error of the mean ( $n = 5$  each); data were analyzed using two-way repeated measures analysis of variance, followed by Tukey's post-hoc test. \*\*\*  $P < 0.001$  vs. colitis group; ###  $P < 0.001$  vs. control group. AV, *A. vulgaris*.

#### 4.3. *A. vulgaris* Amended Serum Levels of MDA, GSH, and SOD Activity in the Rat Model of Colitis

One-way ANOVA revealed a significant difference in serum levels of MDA [ $F(5, 24) = 51.49, P < 0.0001$ ], GSH [ $F(5, 24) = 191.2, P < 0.0001$ ], and SOD activity [ $F(5, 24) = 229.9, P < 0.0001$ ] among the groups. The post-hoc analysis revealed that the colitis rats had significantly increased serum levels of MDA ( $P < 0.001$ ) compared to the control group. However, *A. vulgaris* treatment at 100 and 200 mg/kg significantly decreased MDA levels compared to the colitis group ( $P < 0.001$  to  $P < 0.01$ ) (Figure 3A).

According to Figure 3B, C, a significant reduction in the serum GSH levels and SOD activity was observed in the colitis group compared to the control ( $P < 0.001$ ) (Figure 3B, C). Nevertheless, the administration of AV50, AV100, and AV200 resulted in a significant increase in serum GSH levels and SOD activity in colitis rats compared to untreated colitis rats ( $P < 0.001$ ) (Figure 3B, C). Furthermore, these parameters were significantly higher in the sulfasalazine-treated group compared to the colitis group ( $P < 0.001$ ) (Figure 3B, C).

#### 4.4. *A. vulgaris* Improved Macroscopic Alterations in the Rat Model of Colitis

The results of one-way ANOVA of the macroscopic score indicated a significant difference between the groups [ $F(5, 12) = 19.23, P < 0.0001$ ]. The post-hoc analysis revealed that compared to controls, acetic acid-induced colitis rats displayed severe ulceration and inflammation in the colonic damage ( $P < 0.001$ ) (Figure 4). Meanwhile,

*A. vulgaris*-treated groups showed an amended severity of lesion score compared to the colitis group, with the highest healing effect seen in the groups treated with 200 mg/kg *A. vulgaris* ( $P < 0.01$ ) (Figure 4). Additionally, the rats treated with sulfasalazine had significantly lower macroscopic scores as compared to the colitis group ( $P < 0.001$ ) (Figure 4).


#### 4.5. *A. vulgaris* Alleviated Microscopic Alterations in the Rat Model of Colitis

As depicted in Table 1, the colon of all the animals in the control group showed a typical architecture, including normal colonic mucosa lined with simple columnar epithelium and muscularis mucosae, submucosa with infiltration of a few inflammatory cells, muscularis, and serosa layers (Figure 5A). In contrast, the colon of rats in the groups that received acetic acid showed mild, moderate, and severe tissue changes. The main histopathologic findings in the colon were focal ulceration, necrosis, loss of goblet cells, crypt disarray, submucosal edema, mucosal and submucosal mono- and polymorphonuclear cells infiltration with crypt abscess. The extent and severity of the lesions were more prominent in the acetic acid group (Figure 5B-D). Treatment with the *A. vulgaris* hydroalcoholic extract and sulfasalazine significantly reduced acetic acid-induced pathological lesions in the colon tissues. The administration of sulfasalazine and *A. vulgaris* at a dose of 200 mg/kg showed less focal necrosis, hemorrhage, ulceration, and inflammatory reaction in the mucosa, submucosa, and muscularis layers, reduction in submucosal edema and loss of goblet cell compared to the group treated with *A. vulgaris* at doses of 100 and 50 mg/kg (Figure 5E-H). Regarding the pathological scores, the highest score of colon lesions was observed in the acetic acid control group, followed by *A. vulgaris* extract at 50, 100, and 200 mg/kg and sulfasalazine groups, respectively (Table 1). In the comparison between the groups treated with *A. vulgaris* extract, the most therapeutic effects were observed with the administration of 200 mg/kg of the extract.

## 5. Discussion

This study showed the anti-inflammatory effects of the hydro-alcoholic extract of *A. vulgaris* in experimentally induced acute colitis, characterized by ameliorated colonic lesions, improved body weight, reduced MDA, NO, TNF- $\alpha$  levels, increased SOD activity, and GSH levels in the serum.

Several experimental studies indicated that colitis is triggered by inflammation and oxidative stress (15,



**Figure 2.** Effect of *A. vulgaris* extract treatment on serum levels of nitric oxide (NO) (A); and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) (B) in the acetic acid (AA)-induced colitis rats. Data are presented as mean  $\pm$  standard error of the mean (n = 5 each); data were analyzed using one-way analysis of variance, followed by Tukey's post-hoc test. \*\*\* P < 0.001 vs. colitis group; ### P < 0.001 vs. control group. AV, *A. vulgaris*.


**Table 1.** Histopathological and Inflammatory Response Scores of Colons in Different Groups <sup>a</sup>

Groups	Histopathological Scores		Inflammatory Response Scores	
	Mean $\pm$ Standard Error	Median (Min - Max)	Mean $\pm$ Standard Error	Median (Min - Max)
Control	0.00 $\pm$ 0.00	0 (0 - 1)	0.20 $\pm$ 0.10	0 (0 - 1)
Colitis	11.86 $\pm$ 0.33 <sup>#</sup>	12 (9 - 14)	2.80 $\pm$ 0.10 <sup>#</sup>	3 (2 - 3)
Sulfasalazine	3.93 $\pm$ 0.18 <sup>#</sup>	4 (3 - 5)	1.00 $\pm$ 0.13 <sup>#</sup>	1 (0 - 2)
<i>A. vulgaris</i> (50 mg/kg)	8.60 $\pm$ 0.25 <sup>#</sup>	9 (7 - 10)	2.20 $\pm$ 0.10 <sup>#</sup>	2 (2 - 3)
<i>A. vulgaris</i> (100 mg/kg)	5.93 $\pm$ 0.20 <sup>#</sup>	6 (5 - 8)	1.73 $\pm$ 0.11 <sup>#</sup>	2 (1 - 2)
<i>A. vulgaris</i> (200 mg/kg)	4.40 $\pm$ 0.19 <sup>#</sup>	4 (3 - 6)	1.20 $\pm$ 0.14 <sup>#</sup>	1 (0 - 2)

<sup>a</sup> Data are presented as mean  $\pm$  standard error of the mean (n = 3 each); data were analyzed using one-way analysis of variance, followed by Tukey's post-hoc test <sup>#</sup> vs. control group, <sup>#</sup> vs. colitis group, P < 0.05.

34). Excessive free radical production and decreased antioxidant capacity can affect cellular integrity (35), activation of various transcription factors, and expression of some genes involved in inflammatory pathways that directly and indirectly damage intestinal epithelial cells and lead to cell death (36), demonstrating that flavonoid derivatives in herbs have anti-inflammatory effects, along with marked antioxidant activity and inhibition of enzymes involved in the production of eicosanoids (37, 38). In our study, treatment with *A. vulgaris* (as a potent flavonoid) promoted the serum SOD activity and GSH level, significantly reduced serum TNF- $\alpha$ , NO levels, and lipid peroxidation (MDA), and alleviated colonic damage (39). These results suggest that the anti-inflammatory and antioxidant effects of *A. vulgaris* extract may be due to the presence of flavonoids (40). Few experimental models of UC were performed on other species of the genus

*Artemisia*. In this regard, Shin et al. reported that *Artemisia argyi* extract treatment for 10 days reduced the expression of inflammation-related proteins and genes in the colon and serum, increased antioxidant capacities, and relieved symptoms of UC and body weight in the dextran sodium sulfate (DSS)-induced colitis mouse model (41). Another study showed that treatment of two flavonoid compounds from *Artemisia asiatica* (eupatilin and quercetin) for 4 days before and after inducing colitis in rats reduced the NO production and malondialdehyde levels and increased glutathione levels in the colon. Moreover, this study displayed attenuation in the morphology of the lesions (42). Kalhor et al. demonstrated that *Artemisia dracuncululus* L. administration for 10 consecutive days improved total antioxidant capacity and suppression of pro-inflammatory cytokines in the AA-induced colitis Wistar rat model (43). Furthermore, several studies




**Figure 3.** Effect of *A. vulgaris* extract treatment on serum levels of malondialdehyde (MDA)(A); glutathione (GSH)(B); and superoxide dismutase (SOD)(C) activity in the acetic acid (AA)-induced colitis rats. Data are expressed as mean  $\pm$  standard error of the mean (n = 5 each); data were analyzed using one-way analysis of variance, followed by Tukey's post-hoc test. \*\*P < 0.01, \*\*\*P < 0.001 vs. colitis group; ###P < 0.001 vs. control group. AV, *A. vulgaris*.

reported the beneficial effects of *Artemisia* in models of various inflammatory disorders in the liver (37) and lung (44) tissues. It seems that alleviated damage and improved function of the colon in UC animals by *A. vulgaris* treatment is mediated, at least in part, through the modulation of oxidative stress and inflammatory mediators. Possible mechanisms for this finding are attributed to the inhibition of the expression of genes relevant to the inflammatory process, including COX-2, iNOS, and nuclear factor-kappa B (NF- $\kappa$ B), which reduces the release of pro-inflammatory factors in the tissue (23, 45). Additionally, evidence shows that *Artemisia* extract enhances the expression of enzymes involved in the antioxidant system via activation of the Nrf2 (25).

Thus, the *A. vulgaris* extract can be an effective treatment for colitis. We suggest that future studies assess the confirmatory factors of these changes in the chronic UC model and confirm their subclinical effects in UC patients.

### 5.1 Conclusions

Our findings revealed that *A. vulgaris* extract had protective effects in AA-induced colitis rats in a dose-dependent manner by improving body weight loss and colonic histopathology, decreasing NO and TNF- $\alpha$  levels, and increasing serum antioxidant levels. Thus, it could be considered a potential therapeutic agent against UC.



**Figure 4.** Effect of *A. vulgaris* extract treatment on the microscopic score of colonic injury in the acetic acid (AA)-induced colitis rats. Data were analyzed using one-way analysis of variance, followed by Tukey's post-hoc test. \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$  vs. colitis group; ###  $P < 0.001$  vs. control group. AV, *A. vulgaris*.

## Acknowledgments

The authors would like to express their gratitude to Ms. Mina Salimi for editing the manuscript.

## Footnotes

**Authors' Contribution:** AMJ and MHF contributed to the study conception and design. SS, HK, MH, and SF performed the animal experiments. AMJ and SS analyzed the data and drafted the manuscript. All the authors read and approved the final version.

**Conflict of Interests:** The authors declare no conflict of interest.

**Ethical Approval:** All the procedures were conducted in compliance with the National Institutes of Health guidelines for the care and use of laboratory animals (NIH; publication No. 82-23, revised 1985 and again implemented in 1996) and approved by the Ethics Committee of Ardabil University of Medical Sciences (IR.ARUMS.REC.1400.202).

**Funding/Support:** This work was supported by the Ardabil University of Medical Sciences (grant no. 400000205), Ardabil, Iran.


## References

- Magro F, Gionchetti P, Eliakim R, Ardizzone S, Armuzzi A, Barreiro-de Acosta M, et al. Third European Evidence-based Consensus on Diagnosis and Management of Ulcerative Colitis. Part 1: Definitions, Diagnosis, Extra-intestinal Manifestations, Pregnancy, Cancer Surveillance, Surgery, and Ileo-anal Pouch Disorders. *J Crohns Colitis*. 2017;**11**(6):649-70. [PubMed ID: 28158501]. <https://doi.org/10.1093/ecco-jcc/jjx008>.
- Eisenstein M. Ulcerative colitis: towards remission. *Nature*. 2018;**563**(7730):S33. [PubMed ID: 30405234]. <https://doi.org/10.1038/d41586-018-07276-2>.
- Torres J, Billioud V, Sachar DB, Peyrin-Biroulet L, Colombel JF. Ulcerative colitis as a progressive disease: the forgotten evidence. *Inflamm Bowel Dis*. 2012;**18**(7):1356-63. [PubMed ID: 22162423]. <https://doi.org/10.1002/ibd.22839>.
- Ng SC, Shi HY, Hamidi N, Underwood FE, Tang W, Benchimol EI, et al. Worldwide incidence and prevalence of inflammatory bowel disease in the 21st century: a systematic review of population-based studies. *Lancet*. 2017;**390**(10114):2769-78. [PubMed ID: 29050646]. [https://doi.org/10.1016/S0140-6736\(17\)32448-0](https://doi.org/10.1016/S0140-6736(17)32448-0).
- Shayesteh AA, Saberifirozi M, Abedian S, Sebgatolahi V. Epidemiological, Demographic, and Colonic Extension of Ulcerative Colitis in Iran: A Systematic Review. *Middle East J Dig Dis*. 2013;**5**(1):29-36. [PubMed ID: 24829667]. [PubMed Central ID: PMC3990134].
- Zahedi MJ, Darvish Moghadam S, Hayat Bakhsh Abbasi M, Dehghani M, Shafiei Pour S, Zydabady Nejad H, et al. The incidence rate of inflammatory bowel disease in an urban area of Iran: a developing country. *Middle East J Dig Dis*. 2014;**6**(1):32-6. [PubMed ID: 24829703]. [PubMed Central ID: PMC4005476].
- Hisamatsu T, Kanai T, Mikami Y, Yoneno K, Matsuoka K, Hibi T. Immune aspects of the pathogenesis of inflammatory bowel disease. *Pharmacol Ther*. 2013;**137**(3):283-97. [PubMed ID: 23103332]. <https://doi.org/10.1016/j.pharmthera.2012.10.008>.
- Wu H, Rao Q, Ma GC, Yu XH, Zhang CE, Ma ZJ. Effect of Triptolide on Dextran Sodium Sulfate-Induced Ulcerative Colitis and Gut Microbiota in Mice. *Front Pharmacol*. 2019;**10**:1652. [PubMed ID: 32063856]. [PubMed Central ID: PMC7000629]. <https://doi.org/10.3389/fphar.2019.01652>.
- Yasukawa K, Hirago A, Yamada K, Tun X, Ohkuma K, Utsumi H. In vivo redox imaging of dextran sodium sulfate-induced colitis in mice using Overhauser-enhanced magnetic resonance imaging. *Free Radic Biol Med*. 2019;**136**:1-11. [PubMed ID: 30928473]. <https://doi.org/10.1016/j.freeradbiomed.2019.03.025>.
- Eisenstein M. Biology: A slow-motion epidemic. *Nature*. 2016;**540**(7634):S98-9. [PubMed ID: 28002393]. <https://doi.org/10.1038/540S98a>.
- Rahimi R, Nikfar S, Abdollahi M. Induction of clinical response and remission of inflammatory bowel disease by use of herbal medicines: a meta-analysis. *World J Gastroenterol*. 2013;**19**(34):5738-49. [PubMed ID: 24039370]. [PubMed Central ID: PMC3769914]. <https://doi.org/10.3748/wjg.v19.i34.5738>.
- Wang Z, Li S, Cao Y, Tian X, Zeng R, Liao DF, et al. Oxidative Stress and Carbonyl Lesions in Ulcerative Colitis and Associated Colorectal Cancer. *Oxid Med Cell Longev*. 2016;**2016**:9875298. [PubMed ID: 26823956]. [PubMed Central ID: PMC4707327]. <https://doi.org/10.1155/2016/9875298>.
- Chen S, Wu X, Yu Z. Juglone Suppresses Inflammation and Oxidative Stress in Colitis Mice. *Front Immunol*. 2021;**12**:674341. [PubMed ID: 34421890]. [PubMed Central ID: PMC8375437]. <https://doi.org/10.3389/fimmu.2021.674341>.
- Soufli I, Toumi R, Rafa H, Touil-Boukoffa C. Overview of cytokines and nitric oxide involvement in immuno-pathogenesis of inflammatory bowel diseases. *World J Gastrointest Pharmacol Ther*. 2016;**7**(3):353-60. [PubMed ID: 27602236]. [PubMed Central ID: PMC4986402]. <https://doi.org/10.4292/wjgpt.v7.i3.353>.
- Ansari MN, Rehman NU, Karim A, Soliman GA, Ganaie MA, Raish M, et al. Role of Oxidative Stress and Inflammatory Cytokines (TNF-alpha and IL-6) in Acetic Acid-Induced Ulcerative Colitis in Rats: Ameliorated by *Osteoglyca fruticosa*. *Life (Basel)*. 2021;**11**(3):195.

- [PubMed ID: 33802553]. [PubMed Central ID: PMC8001148]. <https://doi.org/10.3390/life11030195>.
16. Rahimi R, Shams-Ardekani MR, Abdollahi M. A review of the efficacy of traditional Iranian medicine for inflammatory bowel disease. *World J Gastroenterol.* 2010;**16**(36):4504-14. [PubMed ID: 20857519]. [PubMed Central ID: PMC2945480]. <https://doi.org/10.3748/wjg.v16.i36.4504>.
  17. Song M, Li J, Xiong C, Liu H, Liang J. Applying high-resolution melting (HRM) technology to identify five commonly used *Artemisia* species. *Sci Rep.* 2016;**6**:34133. [PubMed ID: 27698485]. [PubMed Central ID: PMC5048426]. <https://doi.org/10.1038/srep34133>.
  18. Ekiert H, Pajor J, Klin P, Rzepiela A, Slesak H, Szopa A. Significance of *Artemisia Vulgaris* L. (Common Mugwort) in the History of Medicine and Its Possible Contemporary Applications Substantiated by Phytochemical and Pharmacological Studies. *Molecules.* 2020;**25**(19):4415. [PubMed ID: 32992959]. [PubMed Central ID: PMC7583039]. <https://doi.org/10.3390/molecules25194415>.
  19. Abiri R, Silva ALM, de Mesquita LSS, de Mesquita JWC, Atabaki N, de Almeida Jr EB, et al. Towards a better understanding of *Artemisia vulgaris*: Botany, phytochemistry, pharmacological and biotechnological potential. *Food Res Int.* 2018;**109**:403-15. [PubMed ID: 29803465]. <https://doi.org/10.1016/j.foodres.2018.03.072>.
  20. Ashok PK, Upadhyaya K. Evaluation of Analgesic and Anti-inflammatory Activities of Aerial Parts of *Artemisia vulgaris* L. in Experimental Animal Models. *J Biol Active Prod Nature.* 2013;**3**(1):101-5. <https://doi.org/10.1080/22311866.2013.782761>.
  21. Bisht D, Kumar D, Kumar R, Dua K, Chellappan DK. Phytochemistry and pharmacological activity of the genus *artemisia*. *Arch Pharm Res.* 2021;**44**(5):439-74. [PubMed ID: 33893998]. [PubMed Central ID: PMC8067791]. <https://doi.org/10.1007/s12272-021-01328-4>.
  22. Nigam M, Atanassova M, Mishra AP, Pezzani R, Devkota HP, Plygun S, et al. Bioactive Compounds and Health Benefits of *Artemisia* Species. *Nat Prod Commun.* 2019;**14**(7). <https://doi.org/10.1177/1934578x19850354>.
  23. Pandey J, Bhusal S, Nepali L, Khatri M, Ramdam R, Barakoti H, et al. Anti-Inflammatory Activity of *Artemisia vulgaris* Leaves, Originating from Three Different Altitudes of Nepal. *ScientificWorldJournal.* 2021;**2021**:6678059. [PubMed ID: 34257625]. [PubMed Central ID: PMC8245213]. <https://doi.org/10.1155/2021/6678059>.
  24. Soon L, Ng PQ, Chellian J, Madheswaran T, Panneerselvam J, Gupta G, et al. Therapeutic potential of *Artemisia vulgaris*: An insight into underlying immunological mechanisms. *J Environ Pathol Toxicol Oncol.* 2019;**38**(3):205-16. [PubMed ID: 31679308]. <https://doi.org/10.1615/JEnvironPatholToxicolOncol.2019029397>.
  25. Park JM, Han YM, Lee JS, Ko KH, Hong SP, Kim EH, et al. Nrf2-mediated mucoprotective and anti-inflammatory actions of *Artemisia* extracts led to attenuate stress related mucosal damages. *J Clin Biochem Nutr.* 2015;**56**(2):132-42. [PubMed ID: 25759519]. [PubMed Central ID: PMC4345182]. <https://doi.org/10.3164/jcbs.14-76>.
  26. Farzaei MH, Khanavi M, Moghaddam G, Dolatshahi F, Rahimi R, Shams-Ardekani MR, et al. Standardization of *Tragopogon graminifolius* DC. Extract Based on Phenolic Compounds and Antioxidant Activity. *J Chem.* 2014;**2014**:425965. <https://doi.org/10.1155/2014/425965>.
  27. Patra R, Padma S, Mukherjee S. An improved method for experimental induction of ulcerative colitis in Sprague Dawley rats. *MethodsX.* 2023;**10**:102158. [PubMed ID: 37091959]. [PubMed Central ID: PMC1013839]. <https://doi.org/10.1016/j.mex.2023.102158>.
  28. Farzaei MH, Ghasemi-Niri SF, Abdolghafari AH, Baeri M, Khanavi M, Navaei-Nigjeh M, et al. Biochemical and histopathological evidence on the beneficial effects of *Tragopogon graminifolius* in TNBS-induced colitis. *Pharm Biol.* 2015;**53**(3):429-36. [PubMed ID: 25471611]. <https://doi.org/10.3109/13880209.2014.923004>.
  29. William Antonio SG, Carmen R SC, Torre Victor E VL, José L CR, Abhel A CP, Cinthya L AV, et al. Hepatoprotective and Nephroprotective Activity of *Artemisia absinthium* L. on Diclofenac-induced Toxicity in Rats. *Pharmacogn J.* 2020;**12**(5):1032-41. <https://doi.org/10.5530/pj.2020.12.146>.
  30. Rakhshandeh H, Heidari A, Pourbagher-Shahri AM, Rashidi R, Forouzanfar F. Hypnotic Effect of *A. absinthium* Hydroalcoholic Extract in Pentobarbital-Treated Mice. *Neurol Res Int.* 2021;**2021**:5521019. [PubMed ID: 33968448]. [PubMed Central ID: PMC8084640]. <https://doi.org/10.1155/2021/5521019>.
  31. Marbun R, Suwarso E, Yuandani. Immunomodulatory Effects of Ethanolic Extract *Artemisia Vulgaris* L. In Male Rats. *Asian J Pharm Clin Res.* 2018;**11**(13):245. <https://doi.org/10.22159/ajpcr.2018.v11i1.26619>.
  32. Farokhi S, Bahrani GR, Babaie A, Farzaei MH. [Protective and Therapeutic Effect of Oleoresin of *Pistacia atlantica* in Acetic Acid-induced Colitis in Rat]. *J Med Plants.* 2019;**4**(72):135-48. Persian. <https://doi.org/10.29252/jmp.4.72.135>.
  33. El-Akabawy G, El-Sherif NM. Zeaxanthin exerts protective effects on acetic acid-induced colitis in rats via modulation of pro-inflammatory cytokines and oxidative stress. *Biomed Pharmacother.* 2019;**111**:841-51. [PubMed ID: 30616083]. <https://doi.org/10.1016/j.biopha.2019.01.001>.
  34. Seril DN, Liao J, Yang GY, Yang CS. Oxidative stress and ulcerative colitis-associated carcinogenesis: studies in humans and animal models. *Carcinogenesis.* 2003;**24**(3):353-62. [PubMed ID: 12663492]. <https://doi.org/10.1093/carcin/24.3.353>.
  35. Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M. Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chem Biol Interact.* 2006;**160**(1):1-40. [PubMed ID: 16430879]. <https://doi.org/10.1016/j.cbi.2005.12.009>.
  36. Kang JY, Kim JM, Park SK, Lee HL, Heo HJ. A Mixture of *Artemisia argyi* and *Saururus chinensis* Improves PM(2.5)-Induced Cognitive Dysfunction by Regulating Oxidative Stress and Inflammatory Response in the Lung and Brain. *Plants (Basel).* 2023;**12**(6):1230. [PubMed ID: 36986919]. [PubMed Central ID: PMC10059966]. <https://doi.org/10.3390/plants12061230>.
  37. Jiang Z, Guo X, Zhang K, Sekaran G, Cao B, Zhao Q, et al. The Essential Oils and Eucalyptol From *Artemisia vulgaris* L. Prevent Acetaminophen-Induced Liver Injury by Activating Nrf2-Keap1 and Enhancing APAP Clearance Through Non-Toxic Metabolic Pathway. *Front Pharmacol.* 2019;**10**:782. [PubMed ID: 31404264]. [PubMed Central ID: PMC6669816]. <https://doi.org/10.3389/fphar.2019.00782>.
  38. Jakovljević MR, Grujičić D, Vukajlović JT, Marković A, Milutinović M, Stanković M, et al. In vitro study of genotoxic and cytotoxic activities of methanol extracts of *Artemisia vulgaris* L. and *Artemisia alba* Turra. *S Afr J Bot.* 2020;**132**:117-26. <https://doi.org/10.1016/j.sajb.2020.04.016>.
  39. Bucchini A, Ricci D, Messina F, Marcotullio MC, Curini M, Giampieri L. Antioxidant and antifungal activity of different extracts obtained from aerial parts of *Inula crithmoides* L. *Nat Prod Res.* 2015;**29**(12):1173-6. [PubMed ID: 25426874]. <https://doi.org/10.1080/14786419.2014.983102>.
  40. Martin DA, Bolling BW. A review of the efficacy of dietary polyphenols in experimental models of inflammatory bowel diseases. *Food Funct.* 2015;**6**(6):1773-86. [PubMed ID: 25986932]. <https://doi.org/10.1039/c5fo00202h>.
  41. Shin JM, Son YJ, Ha IJ, Erdenebileg S, Jung DS, Song DG, et al. *Artemisia argyi* extract alleviates inflammation in a DSS-induced colitis mouse model and enhances immunomodulatory effects in lymphoid tissues. *BMC Complement Med Ther.* 2022;**22**(1):64. [PubMed ID: 35277165]. [PubMed Central ID: PMC8917695]. <https://doi.org/10.1186/s12906-022-03536-x>.
  42. Joo M, Kim HS, Kwon TH, Palikhe A, Zaw TS, Jeong JH, et al. Anti-inflammatory Effects of Flavonoids on TNBS-induced Colitis of Rats. *Korean J Physiol Pharmacol.* 2015;**19**(1):43-50. [PubMed ID: 25605996]. [PubMed Central ID: PMC4297761]. <https://doi.org/10.4196/kjpp.2015.19.1.43>.
  43. Kalhor P, Mohammadzadeh M, Abtahi Froushani SM.



- Anti-inflammatory and antioxidant potential of *Artemisia dracunculus* L. aqueous extract against acetic acid induced ulcerative colitis in male Wistar rats. *Indian J Exp Biol.* 2023;**61**(1):42-50. <https://doi.org/10.56042/ijeb.v6i101.41130>.
44. Mukherjee AA, Kandhare AD, Rojatkar SR, Bodhankar SL. Ameliorative effects of *Artemisia pallens* in a murine model of ovalbumin-induced allergic asthma via modulation of biochemical perturbations. *Biomed Pharmacother.* 2017;**94**:880-9. [PubMed ID: 28810518]. <https://doi.org/10.1016/j.biopha.2017.08.017>.
45. Zamani S, Emami SA, Iranshahi M, Zamani Taghizadeh Rabe S, Mahmoudi M. Sesquiterpene fractions of *Artemisia* plants as potent inhibitors of inducible nitric oxide synthase and cyclooxygenase-2 expression. *Iran J Basic Med Sci.* 2019;**22**(7):774-80. [PubMed ID: 32373299]. [PubMed Central ID: PMC7196345]. <https://doi.org/10.22038/ijbms.2019.34792.8249>.



**Figure 5.** Histopathological lesion in the colon tissue of normal and acetic acid groups (hematoxylin-eosin (H&E) staining). A, normal group, normal architecture of colon tissue (scale bar = 150  $\mu\text{m}$ ); B, acetic acid group, focal ulceration, severe necrosis and crypt destruction in the mucosal layer (scale bar = 150  $\mu\text{m}$ ); C, acetic acid group, severe infiltration of mono- and polymorph nuclear cells in the submucosal layer (scale bar = 70  $\mu\text{m}$ ); D, necrosis of muscle tissue and infiltration of inflammatory cells in the muscularis layer (scale bar = 70  $\mu\text{m}$ ); E, *A. vulgaris* (50 mg/kg) group, moderate to severe crypt destruction and necrosis with moderate inflammation; F, *A. vulgaris* (100 mg/kg) group, moderate crypt destruction and necrosis with moderate inflammation; G, *A. vulgaris* (200 mg/kg) group, mild crypt destruction, and necrosis with mild inflammation; and H, sulfasalazine group, mild crypt destruction with minimal mucosal inflammation.