

85% did not have screening program, 60% endometrial and 5% ovarian cancer .

sexual relation were 67%. 86% of samples did not have any sexual counseling, 57% had at least 3 symptoms of depression, 86% had orgasmic difficulties, 67% had decreased libido and 37% had sexual phobia (due to their appearance in 90%). The average time for sexual education was 5 minutes. All samples believed counseling with their husbands should be helping them .

Discussion: Health worker efforts for saving patient s life are encouraged but proper counseling can promote quality of their life .

Misinformation is barrier to improvement quality, when misinformation is high, we can conclude education and counseling is poor.

Conclusion: couple counseling has a key role in promotion of sexual healthy life, this aspect of education is neglected nowadays.

Email: azarabad74@yahoo.com

32. Possible late effects of pelvic radiotherapy in women

Refahi S¹., Mehnati P. ²

¹Tehran & Ardebil University of Medical Sciences, ²Tabriz University of Medical Sciences

Abstract

While radiotherapy can destroy cancer cells, it can also affect some of the normal cells in the treatment area. It's difficult to predict exactly how your treatment will affect you because radiotherapy affects people in different ways. Radiotherapy can help in the successful treatment of a number of different cancers, including cervical, womb (endometrial), bowel and bladder cancer. It's important to be aware that severe long-term effects of pelvic radiotherapy are uncommon. Radiotherapy is very carefully planned and given to minimise the number of long-term side effects. This study is about the possible side effects and how they can be controlled if they occur.

Keywords: Radiotherapy, late Effect, Women pelvic

Email: soheila52@yahoo.com