

## The 13<sup>th</sup> Iranian and The 2<sup>nd</sup> International Congress of **Microbiology**سیزدهمین کنگره سراسری و



## Antimicrobial Activity of Oleogum Resin Extract & Essential Oil of Ferula gummosa Boiss Against Clinical Strains of Acinetobacter

دومین کنگره بین المللی میکروب شناسی ایران

Fateme Faridafshar<sup>1</sup>; Abbas Ali Imani Fooladi\*<sup>2</sup>; Reza Mirnejad<sup>3</sup>; Alimohammad Asgharian<sup>1</sup>

1-Department of Microbiology, Tonekabon Branch, Islamic Azad University, Tonekabon, Iran 2-Applied Microbiology, Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran 3-Molecular Biology, Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran faridafsharf@yahoo.com

**Background & Objectives:** Acinetobacter spp. is an important cause of nosocomial infections and with widespread resistance to various antibiotics. Ferula gummosa boiss belonging to the family Apiaceae is a wild plant indigenous to Iran. This plant has been used in traditional medicine for its medical properties . phytochemical investigations show F. gummosa boiss is source of biologically active compounds. The aim of this study was to evaluation of the antibacterial activity of oleogum resin extract & essential oil of F. gummosa boiss against Clinical strains of Acinetobacter.

**Methods:** Alcoholic, hydroalcoholic and aqueous extracts from oleogum resin of *Ferula gummosa* were obtained with maceration Methods and Alcoholic, hydroalcoholic and aqueous essential oils was by hydrodistilation methods using a Clevenger apparatus. Solvents evaporated to dryness under reduced pressure in a rotavapor. Antimicrobial activities of extracts and essential oils against of standard and clinical strains of Acinetobacter were evaluated using the agar diffusion methods with disc & well. The minimal inhibitory concentration determined using Broth microdilution methods.

**Results:** Alcoholic essential oil showed higher antibacterial activity while aqueous extract and essential oil don't have considerably antibacterial activity.

**Conclusion:** Alcoholic and hydroalcoholic extracts and essential oils had growth inhibitory effect on the bacteria. Alcoholic essential oil had significantly greater effect on inhibition of growth of Acinetobacter strains.

**Keywords:** Acinetobacter; Antimicrobial Activity; Ferula gummosa