

Phenotype Study of Extended Spectrum β -Lactamase, AmpC and Carbapenemase Producing *E.coli* Obtained Clinical Samples/Babool/Iran

Zahra Shahandeh*; Farahnaz Sadighian; Khadijehbeigom Rekabpour

Medical Laboratory Group, Faculty of Paramedical, Babol University of Medical Sciences, Babol, Iran
shahandeh_z@yahoo.com

Background & Objectives: Antibiotic resistant *E.coli* as the agents of nosocomial and acquired infections is an important therapeutic problem in the world. β -Lactam antibiotic group is used in treatment, commonly. *E.coli* can produce some kinds of Beta lactamase like ESBL, AmpC and carbapenemase against to these antibiotics. Usual lab tests aren't able to detect *E.coli* producing Beta lactamase, then, we decided to detect these bacteria by using of special phenotyping Methods.

Methods: 176 separated *E. coli* from clinical samples were transported to paramedical faculty microbiology lab. *E.coli* species was confirmed by API 20E strip test. Then, ceftazidime and cefotaxime disks (alone and with clavolanic acid), ceftazidime disk (alone and with 3-amino phenylboronic acid) and ceftazidime and 2-Mercaptopropionic acid disks were used for determination of *E. coli* producing ESBL, AmpC and Carbapenemase, respectively. Data was analyzed by SPSS.

Results: Among 176 detected *E.coli*, totally 141(80.2%) B-lactamase producing bacteria and 77 (43.8%), 50 (28.4%) and 14 (8%) bacteria produced ESBL, Carbapenemase and AmpC, respectively. Also, 40 (22.7%) bacteria produced ESBL and carbapenemase enzymes with each other. There was no bacterium that prouduced all kinds of these 3 enzymes.

Conclusion: Regardly to the results, the high percent of Beta-lactamase producing bacteria and bacteria producing two enzymes together are notable. Using of special phenotyping CLSI protocol for determination of these bacteria in the medical labs is suggested.

Keywords: *E. coli*; ESBL; AmpC; Carbapenemase